

**Dayton Area Drug Survey — Table 4**  
**Percent of 7th Graders Reporting Lifetime Use\* (1992-2014)**

<b>Year</b>	<b>1992</b>	<b>1993</b>	<b>1994</b>	<b>1996</b>	<b>1998</b>	<b>2000</b>	<b>2002</b>	<b>2004</b>	<b>2006</b>	<b>2008</b>	<b>2010</b>	<b>2012</b>	<b>2014</b>
	<b>N=2,687</b>	<b>N=3,287</b>	<b>N=4,392</b>	<b>N=1,845</b>	<b>N=2,097</b>	<b>N=2,286</b>	<b>N=2,573</b>	<b>N=3,130</b>	<b>N=3,603</b>	<b>N=3,708</b>	<b>N=2,949</b>	<b>N=3,709</b>	<b>N=2,518</b>
<b>Marijuana</b>	8.8	12.0	16.9	12.6	11.5	10.0	7.5	5.0	4.9	4.0	4.8	5.9	4.7
<b>Crack</b>	1.9	2.4	4.2	3.4	2.5	3.0	1.2	1.4	1.3	1.1	1.1	1.0	0.7
<b>Cocaine HCL</b>	1.6	2.3	4.2	2.9	2.2	2.4	1.2	1.3	1.1	0.8	1.8	0.9	0.8
<b>Inhalants</b>	11.4	14.6	15.1	21.8	15.8	12.1	8.0	9.2	7.3	6.0	7.3	4.5	2.6
<b>Nitrous Oxide</b>	-	-	-	-	-	5.5	2.6	2.5	2.3	2.1	2.7	1.6	1.3
<b>Steroids</b>	1.0	2.3	3.9	0.6	1.0	2.6	1.3	1.1	1.1	0.8	0.8	0.8	0.7
<b>Nitrites</b>	1.2	2.7	4.2	2.9	1.7	-	-	-	-	-	-	-	-
<b>Hallucinogens</b>	2.0	2.7	4.8	3.6	2.4	3.8	1.7	1.5	1.2	1.3	1.6	1.3	0.9
<b>Opiates</b>	1.7	2.9	4.5	2.3	2.1	2.6	1.6	1.6	1.3	1.7	1.5	1.7	0.9
<b>Heroin</b>	-	-	-	-	-	2.3	1.2	1.5	1.0	1.1	1.2	0.9	0.5
<b>Tranquilizers</b>	3.3	3.8	5.2	3.4	3.3	5.7	4.2	4.3	3.3	3.0	3.4	2.8	2.4
<b>Stimulants</b>	4.1	6.0	7.8	7.0	4.2	4.2	1.6	1.5	1.5	1.1	1.1	1.0	0.7
<b>Methamphetamine</b>	-	-	-	-	-	4.0	2.2	1.9	1.4	1.5	1.3	0.9	0.9
<b>Non Rx Diet/SA Pills</b>	-	-	-	-	-	8.9	7.8	6.1	4.4	4.4	1.5	3.6	2.6
<b>PCP</b>	1.8	3.1	4.7	3.6	3.1	-	-	-	-	-	-	-	-
<b>Alcohol</b>	51.1	51.4	51.9	47.8	41.5	38.8	27.3	28.0	23.5	21.2	22.3	17.1	13.5
<b>Cigarettes</b>	32.1	36.0	38.5	36.7	29.3	28.4	19.2	14.0	10.9	8.5	9.5	9.5	6.1
<b>Smokeless Tobacco</b>	11.8	11.7	12.7	10.7	6.9	5.4	3.9	3.7	2.8	2.6	3.0	2.9	2.2
<b>Ritalin</b>	-	-	-	-	-	-	2.6	2.7	1.3	1.0	1.1	0.5	0.6
<b>Ecstasy</b>	-	-	-	-	-	-	1.7	1.6	1.1	0.8	1.2	0.9	0.8

\*Lifetime use -- Having used the drug in question at least once

See: <http://www.med.wright.edu/citar/dads/>